



Rognstranda A beach between two worlds

14

Nome

Siljan

Larda

Skien

Bamb

Porsgrunn

Kragerø





United Nations Educational, Scientific and Cultural Organization Gea Norvegica UNESCO Global Geopark



ROGNSTRANDA – A Geological Divide

Rognstranda – summer paradise and geological phenomenon

At Rognstranda beach you can wander over 1 billion years of geological history in just a few minutes. Two entirely different geological worlds, separated by only a little bay; this is what you meet at Rognstranda.

Just have a little look around and it is obvious! On the one side, there is a relatively flat polished ice-carved bedrock of old gneisses, while on the other side a cliff towers overhead, made of layer upon layer of limestone. The origins of these two types of bedrock are just as dissimilar as the landscapes.

An "unconformity" - what is that?

The gneisses in the foreground of the photo above have a vertical structure. The limestones and shales in the steep cliff in the background lie horizontally. This kind of discontinuous boundary between two rock formations is called an «unconformity», and The glacially-polished bedrock consists of ca. 1500 million year old gneisses from the Precambrian. These rocks were formed deep down in a mountain chain, where they were kneaded and partially melted, before they again cooled and hardened into the metamorphic rock type gneiss. (cont. last page)

at Rognstranda the age difference between the rock layers below and above is over half a billion years. This kind of boundary shows that something geologically very important has occurred. The two sides of this bay belong to two entirely different geological worlds.



On the other side of the bay is the limestone cliff, ca. 500 million years old, from the time periods known as Cambrian, Ordovician and Silurian.

These limestones were formed in a shallow tropical sea when the geological plate that Norway sits on was south of the equator. There was abundant life in the sea, with primitive squids, trilobites, brachiopods, corals, crinoids and bryozoans. Most of these had shells of calcium carbonate, and after they died and sank to the bottom they formed layer upon layer of carbonate ooze, which mixed with clay and sand became compacted into the rock type limestone.

We can of course also find fossils of these extinct organisms in the limestone.



Once sea floor in a shallow sea, now a steep cliff overlooking Rognstranda.

Calcareous pine forest on rocky shelves

This type of natural landscape characterized by limestone cliffs is typical for the Grenland area and is a result of ca. 250 million years of erosion. The cliffs are called «flauer» locally. At the top of several of these «flauer», including the one at Rognstranda, we find protected areas of calcareous pine forest. Calcareous pine forest generally grows where there is limestone bedrock.

Limestone weathers easily and supplies minerals and nutrients to the soil as it disintegrates. Due to this and the acidic carpet of needles that falls from fir trees this area has an unusual vegetation with high species diversity. Among other plants, a rare species of orchid grows on the calcareous forest floor.

A shallow, tropical sea from 400-500 million years ago gave rise to the eastern ridge of Rognstranda.

The photo collage is based on a model by John Klausmeyer, Exhibit Designer, University of Michigan, Museum of Natural History

Contact us at: **post@geanor.no** or visit us at Porselensfabrikken, **Porselensveien 6A, 3920 Porsgrunn**

Like us at facebook

© Gea Norvegica Geopark 2016

Photos taken by the Geopark if otherwise unnamed. Lay-out by J.F.